


High-performance geoscientific computing in multi-scale mineral potential studies



Example of abundant data at different scales from the Mullikkoräme area.


A central task in mine site evaluation and 3D mineral potential studies is to collect data on multiple spatial scales and then use inverse methods to infer the location and extent of economically interesting mineral deposits. Datasets comprise, for example, airborne and ground geophysical data, drill hole data, geological maps and cross sections, drill core logs, and geochemical data. Directly observed geological information is often sparse (e.g. drill holes) and subsurface geology has to be inferred through interpretation and inversion of measured geophysical data.

Project GECCO combines expertise in high performance computing and geomodelling, and aims to develop tools for faster geological modelling in a powerful computing environment.

Common earth model (CEM) is a new concept in multidisciplinary integration of data and work processes. Modern 3D earth modelling software enables sharing common digital 3D representations of the subsurface and rapidly incorporating new information into existing models. 3D models and all geoscientific data can be included in the same CEM model.

Lithological, geochemical and geological properties in the CEM fulfill geological and geophysical constraints. Geological constraints include geological 3D models, geological maps, geochemical data and geological rules inferred from structural geological observations.

The increasing amounts of data has led to the need to use high performance computing techniques in geosciences, with parallel computing and the use of modern accelerator technologies like graphical processing units GPUs to speed up calculations. The first computational results show that GPUs can speed up the direct calculation of Newtonian gravitation by a factor of 215. For instance, a source grid 500x500x40 and a receiver grid 84x84 can be calculated in 5.2 secs. In addition, preliminary results also indicate that the use of the exact formula for the gravitational field of a cube will deviate from that of a spherical source by less than one percent (with a measuring height equal to 1 unit).



Stochastic geophysical inversion requires the computation of a large number of geophysical responses of geological realizations, which is possible with the increased computational capacity.

Researchers

High Performance Computing Group at Åbo Akademi

-Jan Westerholm, PhD, professor, Consortium PI

-Mats Aspñäs, PhD, senior lecturer, post-doctoral researcher

Geological and Geophysical expertise from Geological Survey of Finland:

-Eevaliisa Laine, DSc(Tech), senior scientist, Subproject PI

-Suvi Heinonen, PhD, senior scientist, post-doctoral researcher

-Soile Aatos, MSc, geologist, doctoral researcher

-Kimmo Korhonen, MSc(Tech), geophysicist, doctoral researcher

-Mira Markovaara-Koivisto, MSc(Tech), geologist, doctoral researcher

-Ilkka Suppala, MSc(Tech), geophysicist, researcher

Contact Information

Jan Westerholm <https://research.it.abo.fi/people/jawester>

Åbo Akademi University

Faculty of Science and Engineering

Agora, Vattenborgsvägen

20500 Åbo

+358(0)2 215 4044

jan.westerholm@abo.fi

Eevaliisa Laine

Geological Survey of Finland

Bedrock construction and site investigation unit

Betonimiehenkuja 4

02150 Espoo

+358(0)29 503 2102

eevaliisa.laine@gtk.fi